

The CARE CERTIFICATE

Basic Life Support

/ D V W X S \$ Ø J X H Ø V Y

- What you need to know

Standard

THE CARE CERTIFICATE WORKBOOK

12

Legislation and basic life support



Introduction

The information in standard 12 provides knowledge about the administration of basic life support.

Danger	Response	Airways	Breathing	Call 999/Circulation	Defibrillation
<p>Prior to approaching the casualty, ensure the safety of the casualty, yourself and any bystanders.</p>	<p>If possible, approach the casualty from their feet as this prevents hyperextension of the neck from a responsive casualty. Use the AVPU scale when checking for a response.</p>	<p>Open the airway</p> <p>Place the casualty onto their back. Open the airway using the head-tilt-chin lift method</p> <p>(place your hand on their forehead and gently tilt back the head; with your fingertips under the point of the casualties chin, lift the chin to open the airway)</p>	<p>After opening the airway look, listen and feel for normal breathing for no more than 10 seconds</p>	<p><u>Call an ambulance (999/112)</u> Ask a helper to call otherwise call yourself, stay with the casualty when making the call if possible, activate speaker function on the phone to aid communication with ambulance service.</p>	<p>If an AED arrives, switch it on and follow the spoken or visual prompts. An AED is used in conjunction with CPR.</p>
<p>A – Alert – Is the casualty moving/talking? - No – Proceed to V</p> <p>V – Voice – Do they respond to speech - No – Proceed to P</p> <p>P – PD PHDQLQJ WR FDXVH D switc the ci</p>			<p>Helpful Hint Noisy Gasps In the first few minutes after a cardiac arrest, a casualty may appear to fit and may be barely breathing or taking infrequent, slow noisy gasps. Do not confuse this with normal breathing. If in any doubt that breathing is normal, act as if not breathing normally and prepare to start CPR.</p>	<p>Send someone to get an AED if available and staff are trained to use it. If you're on your own do not leave the casualty. Start CPR</p> <p>Casualty not breathing Commence CPR (30 compressions 2 breaths) Depth of compression 5-6cm at a rate of 100-120 compressions per minute.</p>	

Cardiopulmonary resuscitation (CPR)

Cardiopulmonary resuscitation (CPR) should be administered to a casualty who is not breathing normally and who shows no signs of life. CPR is a method of combining chest compressions with 'effective rescue breaths' in order to artificially circulate blood and to put air into the lungs. The depth of compressions is as follows:

Adult: 5–6 centimetres (similar to the short side of a credit card) using both hands.



Child (1 year to onset of puberty): – compress at least one third of the chest's depth (5cm), using one hand.





Infant (0–1 years of age) – compress at least one third of the chest's depth (4cm), using two fingers.

The rate of compression should be 100–120 compressions per minute. 30 chest compressions should be administered prior to moving on to breaths (called 'expired air ventilation').

After completing 30 chest compressions, two effective breaths should be administered directly into the casualty's mouth, or in the case of an infant, into their mouth and nose. Each breath should take one second to complete and the casualty's chest should rise as in normal breathing; this is known as 'effective rescue breathing'. Turn your head and watch the chest rise and fall, then administer the second breath.

Please note: CPR must be practiced in a simulated environment as part of the Care Certificate training arranged by your employer. The use of this workbook alone is not sufficient to provide you with the skills to perform CPR, and is not sufficient to achieve the competences required for award of the Care Certificate.



Danger	Response	Airways	Breathing	Call 999/Circulation	Defibrillation
<p>Prior to approaching the child or infant, ensure their safety, your safety and the safety of any bystanders.</p>	<p>INFANT</p> <ol style="list-style-type: none"> 1. Talk to the infant. 2. Gently stimulate the infant. 3. If a response is gained, check for further injuries (secondary survey) and contact the emergency services if required. <p>CHILD</p> <ol style="list-style-type: none"> 1. Talk to the child 2. Gently stimulate the child and ask loudly 'Are you alright?' 3. If a response is gained, check for further injuries (secondary survey) and contact the emergency services if required. 	<p>Open the airway</p> <p>Place the infant/child onto their back. Open the airway using the head-tilt-chin lift method</p> <p>(place your hand on their forehead and gently tilt back the head; with your fingertips under the point of the infants/child's chin, lift the chin to open the airway)</p>	<p>After opening the airway look, listen and feel for normal breathing for no more than 10 seconds</p> 	<p><u>Call an ambulance (999/112)</u> Ask a helper to call otherwise call yourself. If you are on your own perform CPR for 1 min before going for help (5 initial rescue breaths before starting chest compressions). Stay with the casualty when making the call if possible or if able to carry the infant or child whilst summoning help, activate speaker function on the phone to aid communication with ambulance service. <u>Send someone to get an AED if available</u></p> <p>Casualty not breathing Commence CPR, 5 initial rescue breaths (30 compressions 2 breaths) Depth of compression 4 cm for an infant, 5 cm for a child at a rate of 100-120 compressions per minute.</p>	<p>If an AED arrives, switch it on and follow the spoken or visual prompts. An AED is used in conjunction with CPR.</p>
					

Obstructed airway (adult)

The main aim of the respiratory system is to supply oxygen to all parts of the body. Breathing is essential to life.

The airway can be obstructed in a variety of ways including foreign bodies (food or other items), allergic reactions, asthma, blood, vomit and infections. An obstruction can cause minor or major breathing difficulties and, in severe circumstances, may cause the casualty to become unconscious and unresponsive.

Someone who is choking will have either a partial or complete obstruction of the airway. The severity of the blockage will determine the difficulty in breathing.

Recognition

- Grasping at the throat area.
- Difficulty in breathing and speaking.
- Difficulty in crying or making a noise.
- Redness of the face.
- Eyes enlarged and watering.
- Displaying distress.

Treatment

This should not be carried out by any worker who has not successfully completed formal practical training provided by their employer.
Encourage the casualty to lean forward and cough.

If the obstruction remains:

administer a maximum of 5 sharp blows between the shoulder blades.

If the obstruction still remains:

administer a maximum of 5 abdominal thrusts (or chest thrusts for an infant).

, I W K H R E V W U X F W L R Q L V V W L O O Q R W U H O L H Y H G F D O O
E D F N E O R Z V Z L W K I L Y H D E G R P L Q D O W K U X V W V

Start CPR if the casualty becomes unresponsive:

- Support the casualty carefully to the ground.
- Immediately call the ambulance service.
- Begin CPR with chest compressions.

Obstructed airway (infant and child)

An obstruction can cause minor or major breathing difficulties and, in severe circumstances, may cause the infant or child to become unconscious or unresponsive.

Recognising a choking infant or child

- Grasping at the throat area.
- Difficulty in breathing and speaking (in the case of a child).
- Difficulty in crying or making a noise.
- Redness of the face.
- Eyes enlarged and watering.
- Displaying distress.

With a complete obstruction the infant or child may show the above signs but also the skin colour may develop a blue/grey tinge; they will get progressively weaker and eventually they will become unconscious.



Treating a choking infant

Consider the safest action to manage the choking child:

If the infant is coughing effectively, then no external manoeuvre is necessary, monitor continuously. If the infant's coughing is, or is becoming, ineffective, shout for help immediately and determine the infant's conscious level.

Conscious choking infant

If the infant is still conscious but has absent or ineffective coughing, give back blows.

If back blows do not relieve choking, give chest thrusts. These manoeuvres create an 'artificial cough' to increase intrathoracic pressure and dislodge the foreign body.

Back blows

Support the infant in a head-downwards, prone position, to enable gravity to assist removal of a foreign body, a seated or kneeling first aider should be able to support the infant safely across their lap.

Support the infant's head by placing the thumb of one hand at the angle of the lower jaw, and one or two fingers from the same hand at the same point on the other side of the jaw.

Do not compress the soft tissues under the infant's jaw, as this will exacerbate the airway obstruction.

Deliver up to 5 sharp back blows with the heel of one hand in the middle of the back between the shoulder blades.

The aim is to relieve the obstruction with each blow rather than to give all 5.

Chest thrusts

Turn the infant into a head-downwards supine position. This is achieved safely by placing your free arm along the infant's back and encircling the occiput with your hand.

Support the infant down your arm, which is placed down (or across) your thigh. Identify the landmark for chest compression (lower sternum approximately a finger's breadth above the xiphisternum i.e. lowest part of the breastbone).

Deliver up to 5 chest thrusts. These are similar to chest compressions, but sharper in nature and delivered at a slower rate.

The aim is to relieve the obstruction with each thrust rather than to give all 5.

Following chest thrusts reassess the infant:

If the object has not been expelled and the infant is still conscious, continue the sequence of back blows and chest thrusts.

Call out, or send, for help if it is still not available.

Do not leave the child at this stage.

If the object is expelled successfully, assess the infant's clinical condition. It is possible that part of the object may remain in the respiratory tract and cause complications. If there is any doubt, seek medical assistance.

Helpful Hint

Under no circumstances should abdominal thrusts be performed on an infant. These must be replaced with chest thrusts.

Treating a choking child

Consider the safest action to manage the choking child:

If the infant is coughing effectively, then no external manoeuvre is necessary, monitor continuously.

If the infants' coughing is, or is becoming, ineffective, shout for help immediately and determine the infant's conscious level.

Conscious choking child

If the child is still conscious but has absent or ineffective coughing, give back blows. If back blows do not relieve choking, give abdominal thrusts. These manoeuvres create an 'artificial cough' to increase intrathoracic pressure and dislodge the foreign body.

Back blows (In a child over 1 year)

Back blows are more effective if the child is positioned head down.

A small child may be placed across the rescuer's lap as with an infant.

If this is not possible, support the child in a forward-leaning position and deliver the back blows from behind.

Give 5 sharp blows between the shoulder blades with the heel of the other hand.

If back blows fail to dislodge the object, and the child is still conscious, use abdominal thrusts.

Abdominal thrusts for children over 1 year:

Stand or kneel behind the child. Place your arms under the child's arms and encircle their torso.

Clench your fist and place it between the umbilicus (naval) and the ribcage.

Grasp this hand with your other hand and pull sharply inwards and upwards.

Repeat up to 4 more times.

If the obstruction is still not relieved continue alternating 5 back blows with 5 abdominal thrusts.

Start CPR if the child becomes unresponsive

Support the child carefully to the ground.

Immediately call the ambulance service.

Begin CPR with chest compressions.

Confidentiality

Anyone who is responsible for the storage of records and information must be aware of their responsibilities under the General Data Protection Regulation 2016 and, if relevant, the Freedom of Information Act (FOI) 2000.

Record keeping

Depending on your specific job role, there will be information and records that will require completion should an infant, child or adult be involved in an accident, or become ill whilst in the health or social care setting.

This recorded information in the accident book can:

- help to identify trends

- help to control health and safety risks

- be used for reference in future first aid needs assessments

- prove useful for investigations.

Please refer to your employer's policy and procedures and forms. If you need further clarity, speak with your manager.

Refresher training

Refresher training should be conducted regularly. It is good practice to complete a refresher session on basic life support annually. This is not a condition of completion for the Care Certificate.

The CARE CERTIFICATE

Basic Life Support

- What do you know now?

Standard

12

Activity 12.1a



Basic life support

Complete the following sentence with the missing words.

knowledge

competently

practical

practice

employer

You should be provided with training by your
to be able to put this into and in order to be
able to carry out basic life support

Activity 12.2

This section contains five horizontal blue bars, stacked vertically, intended for the student to write their responses to the activity.

Activity 12.3



Link the word on the left to the correct description on the right.

Activity 12.4



Answer the question by filling in the blanks using the numbers provided.

Activity 12.5

Regarding a choking casualty (adult): please place the following in order of action by labelling them 1–4.



Change log

Date of change	Location in document	Original content	Amended content
September 2021	Page 6, Treatment section, Fifth bullet point.	If the obstruction is still not relieved, continue alternating five back blows with five abdominal thrusts and call for help.	If the obstruction is still not relieved, call for help and continue alternating five back blows with five abdominal thrusts.
August 2024	Page 3, reference to APVU	P – Place, place your hand on their shoulders and gently shake them...	P – Pain, meaning to cause a minor pain to see if the person responds. Examples include pinching the ear lobes or finger tips.